

Durham Research Online

Deposited in DRO:

12 January 2017

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Gröcke, D. R. and Gillikin, D. P. (2008) 'Advances in mollusc sclerochronology and sclerochemistry : tools for understanding climate and environment.', *Geo-marine letters.*, 28 (5-6). pp. 265-268.

Further information on publisher's website:

<http://dx.doi.org/10.1007/s00367-008-0108-4>

Publisher's copyright statement:

The final publication is available at Springer via <https://doi.org/10.1007/s00367-008-0108-4>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Advances in mollusc sclerochronology and sclerochemistry: tools for understanding climate and environment

Darren R. Gröcke¹ & David P. Gillikin²

1. Department of Earth Sciences, Durham University, Science Laboratories, Durham DH1 3LE, UK

2. Department of Earth Science & Geography, Vassar College, Box 475, Poughkeepsie, NY 12604, USA

Abstract This Special Issue compiles papers on marine, estuarine and freshwater mollusc shells as recorders of environmental and climatic conditions. Considering that many past studies have differentiated geochemical investigations from sclerochronological investigations, we propose that the sub-discipline term “sclerochemistry” be used when geochemical investigations are undertaken. This issue starts with two review papers that discuss the importance of physiology or vital effects on both sclerochronology and sclerochemistry. Several sclerochemical calibration studies on modern specimens of both bivalves and gastropods are presented (including $\delta^{18}\text{O}$, Mg/Ca, Sr/Ca and Ba/Ca), which illustrate the usefulness and difficulties associated with using these proxies. Studies on fossil mollusc shells are also provided, with one study that uses Pliocene scallop shells to understand past ocean circulation and another that addresses the problem of diagenesis. Finally, a sclerochronological study of crystal prism width across the shell is presented. This issue demonstrates that many elemental and isotopic proxies contained in mollusc shells are complex. In spite of these complexities, environmental and climatic conditions can be extracted from them for use in palaeoclimatic and palaeoenvironmental research.

Introduction

With future climate change at the forefront of environmental policy making (IPCC 2007), studies of past climatic changes and their environmental effects are vital because they allow an understanding of the processes responsible for these changes. By using the past as the key to the future, we can better predict how the Earth will respond to certain environmental perturbations. Information about past climatic conditions and changes can be obtained through proxies; these are geochemical or physical signals recorded in biological or geological structures that reflect the environment in which they formed. Because they typically accrete or deposit sequentially through time they can thus record a time-series of environmental information. For example, environmental information has been obtained from tree rings, sclerosponges, speleothems, corals, fish otoliths, foraminifera, mammal teeth, laminated sediments, ice cores and mollusc shells.

The term sclerochronology was first applied to a radiographic study on corals by Buddemier et al. (1974, p. 196) and more formally by Hudson et al. (1976) in their paper on corals entitled,

“Sclerochronology: a tool for interpreting past environments.” However, more recently at the 1st International Sclerochronology Conference, Jones et al. (2007) define sclerochronology as:

“the study of physical and chemical variations in the accretionary hard tissues of organisms, and the temporal context in which they formed. Sclerochronology focuses primarily upon growth patterns reflecting annual, monthly, fortnightly, tidal, daily, and sub-daily increments of time entrained by a host of environmental and astronomical pacemakers. Familiar examples include daily banding in reef coral skeletons or annual growth rings in mollusk shells. Sclerochronology is analogous to dendrochronology, the study of annual rings in trees, and equally seeks to deduce organismal life history traits as well as to reconstruct records of environmental and climatic change through space and time.”

Many authors differentiate sclerochronological studies and geochemical studies, for example, Williams et al. (1982) state that they used “coordinated isotopic *and* sclerochronological (growth increment) studies” and Schöne et al. (2007) entitled their paper “Combined sclerochronologic and oxygen isotope analysis of gastropod shells...”. Therefore, in line with the differentiation already used by dendrochronologists to separate studies of tree ring width increments (dendrochronology) from isotopic or chemical analyses across rings (dendrochemistry) (e.g., Smith and Shortle 1996; Verheyden et al. 2005; Poussart et al. 2006), we propose to use **sclerochronology** for studies of the physical structure of the hard tissues of organisms, even when combined with geochemistry (e.g., growth line periodicity) and that **sclerochemistry**, as a sub-discipline of sclerochronology, be used to describe solely geochemical (isotopic or elemental) studies of the hard tissues of organisms. Whether these terms and distinctions are adopted or not, it does provide an essential difference between the two approaches.

For more than 50 years, mollusc shells have been known archives of past environmental conditions (Davenport 1938; Epstein et al. 1953; Emiliani et al. 1964; Clark 1968). During mollusc growth they sequentially deposit new layers of shell, and the physical structure and/or chemical composition of these layers may reflect the environmental conditions at the time they formed. Indeed, environmental data have been extracted from both mollusc shell geochemistry (sclerochemistry), as well as from external or internal growth increments (sclerochronology). Molluscs are beneficial in that they can provide high-resolution seasonal records of environmental conditions and have a wide geographic distribution, whereas many other archives, such as scleractinian corals are limited in their latitudinal extent. Although most molluscs typically live less than 10 years, some readily achieve 50 years (Peterson 1986) and there have been reports of bivalves (*Arctica islandica*) living up to 374 years (Schöne et al. 2005) or even older. In addition, bivalve shells are often found in archaeological middens or as fossils, potentially allowing records of environmental conditions to be extended not only into historical records but geologic time.

However, it is becoming increasingly clear that the animals' physiology significantly impacts proxies recorded in the shell.

In 2005 a special issue on environmental records in accretionary hard parts of aquatic organisms including bivalved molluscs, fish, corals and coralline sponges was published (Schöne and Surge 2005). These papers have renewed the interest in the use of molluscs as recorders of past environmental conditions, which in turn led to workshops and conferences in 2007, such as a PAGES and SSHRC funded workshop, "Stable Isotopes in Archaeological Midden Shells: High-Resolution Paleoclimatic & Paleoenvironmental Archives" held between 10–13 July at the Parks Canada Discovery Centre in Hamilton, Canada (Fig. 1), and the 1st International Sclerochronology Conference between 17–21 July in St. Petersburg, Florida. During the discussions at the PAGES/SSHRC workshop participants voted to initiate a Sclerochronology Listserv, which David Goodwin at Denison University volunteered to set up and administer (<https://listserv.cc.denison.edu/www/info/sclerochronology>). The papers within this Special Issue derive mainly from the PAGES/SSHRC meeting and an open invitation through the Sclerochronology Listserv, and focus on mollusc shell sclerochronology and sclerochemistry (both elemental and isotopic studies) in marine, estuarine and freshwater molluscs.

To set the scene, two review papers that cover the topic of physiology are provided. **Schöne** provides an overall view of how physiological effects can affect a range of geochemical proxies in molluscs. By taking a modeling approach with published and unpublished data he looks at investigating periods of growth cessation, how long are they and what is influencing them. Subsequently by looking at variable growth rates a greater understanding of the geochemical proxies can be obtained, thus eliminating these effects in our sclerochemical records.

McConnaughey and Gillikin, however deal specifically with carbon-isotope ratios in molluscs and how metabolic carbon can affect their distribution in the shell. With a process orientated model they determine the percentage of metabolically respired carbon versus dissolved inorganic carbon taken up into the shell. Very few isotope sclerochemical studies report carbon isotopes but instead only publish the oxygen isotope ratios. With this model and understanding of carbon isotopes in shells it should be possible to determine ecosystem metabolism versus estuarine influences on shell geochemistry.

Two modern-based coordinated sclerochronological and sclerochemical studies follow, which are extremely important if one is to use archaeological and/or fossil mollusc shells to interpret environment and climate. **Andrus and Rich** conduct an oxygen-isotope study of *Rangia cuneata*, which inhabits an estuarine setting. Using oxygen isotopes, time-series seawater temperature data and various environmental assumptions they produce a model–data comparison, which indicates that the microstructural increments of this species are controlled by tidal rhythms.

Mannino et al. apply a similar approach but on the gastropod *Osilinus turbinatus* from the

Mediterranean region. Several populations of the gastropod were analyzed for oxygen isotopes and compared against measured sea surface temperatures. Growth lines however occur during the summer when temperatures reach a maximum. They report that the oxygen isotopes are recording the temperature all-year round and have minimal offsets, thus can be used as a good climate indicator from archaeological sites.

Goewert and Surge have adopted the above approach by analyzing the Pliocene scallop, *Chesapecten madisonius*, from Virginia in order to determine seasonality. By using the presence of growth lines as markers of a season they use oxygen isotopes to determine if they occurred in winter or summer. The results indicate that the growth lines occur during winter and not summer, which is common for molluscs from cold-temperate regions, but not warm-temperate faunas. They conclude that this may be the result of a change in the Labrador Current and/or the migration of warm-water temperate species following the Gulf Stream and thus during cold intervals it resulted in a reduction of shell growth. **Lu** uses an alternative approach to determine whether fossil mollusc shells preserve a pristine isotopic signature that can be used for palaeoenvironment and palaeoclimate reconstructions. The Miocene shells in Lu's study were evaluated for an array of elemental and isotopic signals to determine the level of diagenetic alteration, if any. Using modern seawater geochemical baselines it was found that although these Miocene shells look pristine because their fabric had been maintained, they had undergone diagenetic alteration. This type of approach suggests that all future deep-time sclerochemical studies should incorporate a suite of geochemical investigations before using the data to reconstruct palaeo-seawater conditions.

The next two studies assess the potential of trace element proxies in estuarine bivalves. **Gillikin et al.** investigate the barium signal in calcite and aragonite bivalve shells (*Pecten maximus*, *Saxidomus giganteus*). They have previously shown that the background Ba/Ca signal is related to water Ba/Ca ratios in *Mytilus edulis* shells, but here focus on the episodic nature of the Ba/Ca peaks. The Ba/Ca peaks are highly reproducible between specimens, and do not seem to be related to phytoplankton blooms as has been previously proposed. Other possible causes such as dissolved Ba in ambient water, spawning, shell organic matter content and kinetic growth rate effects are discussed, but none provide satisfactory explanations for these Ba/Ca peaks. However, the background signal does seem to be related to water Ba concentrations in these species providing further evidence that this is a useful proxy. **Wanamaker et al.** cultured juvenile mussels (*Mytilus edulis*) in different temperatures and salinities and found that the shell Mg/Ca and Sr/Ca ratios were not well correlated with water temperature. However, when using shells grown in the lowest salinity, the relationships between shell elemental chemistry and water temperature improve moderately ($R^2 = 0.75$ and 0.82 , for Sr/Ca and Mg/Ca respectively). This study highlights the difficulty with these proxies, but does give hope that they may be useful in some environmental settings.

The last two studies in this issue deal with freshwater bivalves. **Carroll and Romanek** studied the sclerochemistry of five specimens of *Elliptio complanata* from four streams (i.e., 20 shells). They compared Mn, Cu, Sr and Ba of the inner and outer shell layers and found that both record accurate environmental information. When comparing shell elemental concentrations to water concentrations they found a strong correlation for Mn, Sr and Ba, but only after excluding five shells from a polluted stream. Their results suggest that sclerochemistry of shells from polluted streams do not accurately record stream geochemistry. This is one of the first studies to clearly illustrate that elemental concentrations in freshwater mussel shells can be used to obtain environmental information. Finally, **Vancolen and Verrecchia** investigated detailed sclerochronology of *Unio tumidus* shells. Their study focused on the distribution of the prism width inside the prismatic layer of the shell, which has implications for growth-line formation and proxy incorporation. They found that prism widths are randomly distributed and are not related to either growth lines or environmental parameters. Crystal morphology has been noted to be responsible for trace element incorporation in inorganic calcite, therefore this study should be repeated on several species of molluscs to ascertain the influence of crystal morphology on sclerochemistry.

The present volume would not have been possible without the thorough and timely work of the reviewers. We express our gratitude to the following (in alphabetical order): Fred Andrus, Karem Azmy, George Clark II, Yannicke Dauphin, Frank Dehairs, David Dettman, Laurent Emmanuel, Julie Ferguson, Pedro Freitas, Ann Goewert, David Goodwin, Dorothee Hippler, Doug Jones, Stuart Jones, David Krantz, Claire Lazareth, Feng Lu, Ted McConnaughey, William Patterson, Yves-Marie Paulet, Chris Romanek, Bernd Schöne, Donna Surge, Maurice Tucker, Alan Wanamaker, and two reviewers who wished to remain anonymous in this list. We also, of course, thank all of the authors who contributed their stimulating research to this issue. Furthermore, we thank the editors Burg Flemming and Monique Delafontaine for making this issue a realization. Thanks to PAGES and SSHRC who provided funding for the workshop at the Parks Canada Discovery Centre in Canada – it was a truly wonderful experience for all of us. In conclusion, we hope that this Special Issue will interest many readers and generate new research in the topics of sclerochronology and sclerochemistry.

Figure 1. Participants of the PAGES and SSHRC funded workshop, “Stable Isotopes in Archaeological Midden Shells: High-Resolution Paleoclimatic & Paleoenvironmental Archives”, 10–13 July, Parks Canada Discovery Centre, Hamilton, Canada.

References

- Buddemier RW, Maragos JE, Knutson DW (1974) Radiographic studies of reef coral exoskeletons: rates and patterns of coral growth. *J Exp Mar Biol Ecol* 14:179-200
- Clark II GR (1968) Mollusk shell: Daily growth lines. *Science* 161:800-802
- Davenport CB (1938) Growth lines in fossil pectens as indicators of past climates. *J Paleontol* 12:514-515
- Emiliani C, Cardini L, Mayeda T, McBurney CBM, Tongiorgi E (1964) Palaeotemperature analysis of marine molluscs (food refuse) from the site of Arene Candide Cave, Italy and the Haua Fteah Cave, Cyrenaica. In Craig H, Miller SL, Wasserburg GJ (eds), *Isotopic and Cosmic Chemistry*. North Holland Publishing Company, Amsterdam, pp. 133-156
- Epstein S, Buchsbaum R, Lowenstam HA, Urey HC (1953) Revised carbonate - water isotopic temperature scale. *Bull Geol Soc Am* 64:1315-1326
- Hudson JH, EA Shinn, RB Halley, Lidz B (1976) Sclerochronology: a tool for interpreting past environments. *Geology* 4:361-364
- IPCC (2007) *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds.), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp
- Jones DS, Arnold B, Quitmyer I, Schöne, BR, Surge D (2007) 1st International Sclerochronology Conference. <http://conference.ifas.ufl.edu/sclerochronology/> accessed 28 April 2008
- Peterson CH (1986) Quantitative allometry of gamete production by *Mercenaria mercenaria* into old-age. *Mar Ecol-Prog Ser* 29:93-97
- Poussart PM, Myneni SCB, Lanzirrotti A (2006) Tropical dendrochemistry: A novel approach to estimate age and growth from ringless trees. *Geophys Res Lett* 33, L17711, doi:10.1029/2006GL026929
- Schöne BR, Surge D (2005, editors) *Looking back over skeletal diaries — High-resolution environmental reconstructions from accretionary hard parts of aquatic organisms*. *Palaeogeogr Palaeoclimatol Palaeoecol* 228:1-191
- Schöne BR, Fiebig J, Pfeiffer M, Gleß R, Hickson J, Johnson ALA, Dreyer W, Oschmann W (2005) Climate records from a bivalved Methuselah (*Arctica islandica*, Mollusca; Iceland). *Palaeogeogr Palaeoclimatol Palaeoecol* 228:130-148
- Schöne BR, Rodland DL, Wehrmann A, Heidel B, Oschmann W, Zhang Z, Fiebig J, Beck L (2007) Combined sclerochronologic and oxygen isotope analysis of gastropod shells (*Gibbula cineraria*, North Sea): life-history traits and utility as a high-resolution environmental archive for kelp forests. *Mari Bio* 150:1237-1252
- Smith KT, Shortle WC (1996) Tree biology and dendrochemistry. In: Dean JS, Meko DM, Swetnam TW (eds), *Tree Rings, Environment and Humanity*. Radiocarbon, Tucson, pp. 629-635
- Verheyden A., Roggeman M, Bouillon S, Elskens M, Beeckman H, Koedam N (2005) Comparison between $\delta^{13}\text{C}$ of α -cellulose and bulk wood in the mangrove tree *Rhizophora mucronata*: implications for dendrochemistry. *Chem Geol* 219:275-282
- Williams DF, Arthur MA, Jones DS, Healy-Williams N (1982) Seasonality and mean annual sea surface temperatures from isotopic and sclerochronological records. *Nature* 296:432-434

Manuscript order, contacting authors during publication process (*) & received/accepted dates

Gröcke* & Gillikin*: Advances in mollusc sclerochronology and sclerochemistry: a tool for understanding climate and environment

d.r.grocke@durham.ac.uk, dagillikin@vassar.edu

Schöne*: The curse of physiology – Challenges and opportunities in the interpretation of geochemical data from mollusk shells

schoeneb@uni-mainz.de

Received: 20 October 2007

Accepted: 20 February 2008

McConnaughey* & Gillikin*: Carbon isotopes in mollusk shell carbonates

mcconnat@gmail.com, dagillikin@vassar.edu

Received: 20 October 2007

Accepted: 18 April 2008

Andrus* & Rich; A preliminary assessment of oxygen isotope fractionation and growth increment periodicity in the estuarine clam *Rangia cuneata*

fandrus@geo.ua.edu

Received: 24 October 2007

Accepted: 4 April 2008

Mannino*, Thomas*, Leng, Sloane; Shell growth and oxygen isotopes in the topshell *Osilinus turbinatus*: resolving past inshore sea surface temperatures

tcrnmam@ucl.ac.uk, tcfa312@ucl.ac.uk

Received: 30 October 2007

Accepted: 27 February 2008

Goewert* & Surge*; Seasonality and growth patterns using isotope sclerochronology in shells of the Pliocene Scallop, *Chesapecten madisonius*

annieg@unc.edu, donna64@unc.edu

Received: 15 November 2007

Accepted: 12 March 2008

Lu*; Pristine or altered: Low-Mg calcite shells survived from massive dolomitization? A case study from the Miocene carbonates

Lu.Feng@epamail.epa.gov

Received: 30 October 2007

Accepted: 20 February 2008

Gillikin*, Lorrain, Paulet, André, Dehairs; Synchronous barium peaks in high-resolution profiles of calcite and aragonite marine bivalve shells

dagillikin@vassar.edu

Received: 12 November 2007

Accepted: 15 March 2008

Wanamaker Jr.*, Kreutz, Wilson, Borns Jr., Introne, Feindel; Experimentally determined Mg/Ca and Sr/Ca ratios in juvenile bivalve calcite for *Mytilus edulis*: Implications for paleotemperature reconstructions

a.wanamaker@bangor.ac.uk

Received: 25 October 2007

Accepted: 15 February 2008

Carrol & Romanek*; Shell layer variation in trace element concentration for the freshwater bivalve, *Elliptio complanata*

romanek@uga.edu

Received: 2 November 2007

Accepted: 7 April 2008

Vancolen* & Verrecchia*; Does prism width from the shell prismatic layer have a random distribution?

severine.vancolen@unine.ch, eric.verrecchia@unine.ch

Received: 24 October 2007

Accepted: 4 March 2008